Decomposition of optical force into conservative and nonconservative components

Yikun Jiang[1,2], Jun Chen[3], Jack Ng[4], and Zhifang Lin[1,2,5]

[1] *State Key Laboratory of Surface Physics and Department of Physics, Fudan University, Shanghai 200433, China*

[2] *Key Laboratory of Micro and Nano Photonic Structures (Ministry of Education), Fudan University, Shanghai 200433, China*

[3] *Institute of Theoretical Physics and Collaborative Innovation Center of Extreme Optics, Shanxi University, Shanxi, China*

[4] *Department of Physics and Institute of Computational and Theoretical Studies, Hong Kong Baptist University, Hong Kong*

[5] *Collaborative Innovation Center of Advanced Microstructures, Fudan University, Shanghai 200433, China*

arXiv:1604.05138v1 [physics.optics] 18 Apr 2016

We present a multipole expansion theory for optical force exerting on a particle immersed in generic monochromatic free-space optical field. Based on the theory, we have, for the first time, successfully decomposed the optical force on a spherical particle of arbitrary size into a conservative and a nonconservative parts, which are, respectively, written as a gradient of a scalar function and curl of a vector function in an explicit and analytical form. As a result, a scalar potential and a vector potential can be defined, up to gauge freedoms, for the optical force. The decomposition shed light on the understanding of the optical force and pave a new way to engineer optical force for various purposes such as equilibrium statistical mechanics as well as optical micromanipulation.

PACS numbers: 42.50.Tx, 42.50.Wk, 87.80.Cc, 78.70.-g

I. MULTIPOLE EXPANSION OF OPTICAL FORCE

Physically, the period-averaged optical force acting on a particle in any monochromatic optical field is written in two terms, the interception (extinction) force $\langle\vec{\boldsymbol{F}}_{\rm int}\rangle$ and recoil force $\langle\vec{\boldsymbol{F}}_{\rm rec}\rangle$. Based on the T-matrix method [1, 2], the multipole field theory [3, 4], and the irreducible tensor theory [5], the optical force can be written in terms of the electric and magnetic multipoles of various orders induced on the particle. To be specific, the interception force $\langle\vec{\boldsymbol{F}}_{\rm int}\rangle$ is given by

$$\langle\vec{\boldsymbol{F}}_{\rm int}\rangle = \sum_{l=1}^{\infty}\langle\vec{\boldsymbol{F}}_{\rm int}^{(l)}\rangle, \tag{1a}$$

$$\langle\vec{\boldsymbol{F}}_{\rm int}^{(l)}\rangle = \langle\vec{\boldsymbol{F}}_{\rm int}^{{\rm e}(l)}\rangle + \langle\vec{\boldsymbol{F}}_{\rm int}^{{\rm m}(l)}\rangle \tag{1b}$$

$$\langle\vec{\boldsymbol{F}}_{\rm int}^{{\rm e}(l)}\rangle = \frac{1}{2\,l!}\,{\rm Re}\Big[(\nabla^{(l)}\vec{\boldsymbol{E}}^{*})\overset{(l)}{\cdot\cdot}\overleftrightarrow{\mathbb{O}}_{\rm elec}^{(l)}\Big] \tag{1c}$$

$$\langle\vec{\boldsymbol{F}}_{\rm int}^{{\rm m}(l)}\rangle = \frac{1}{2\,l!}\,{\rm Re}\Big[(\nabla^{(l)}\vec{\boldsymbol{B}}^{*})\overset{(l)}{\cdot\cdot}\overleftrightarrow{\mathbb{O}}_{\rm mag}^{(l)}\Big] \tag{1d}$$

and the recoil force $\langle\vec{\boldsymbol{F}}_{\rm rec}\rangle$ reads

$$\langle\vec{\boldsymbol{F}}_{\rm rec}\rangle = \sum_{l=1}^{\infty}\langle\vec{\boldsymbol{F}}_{\rm rec}^{(l)}\rangle, \tag{2a}$$

$$\langle\vec{\boldsymbol{F}}_{\rm rec}^{(l)}\rangle = \langle\vec{\boldsymbol{F}}_{\rm rec}^{{\rm e}(l)}\rangle + \langle\vec{\boldsymbol{F}}_{\rm rec}^{{\rm m}(l)}\rangle + \langle\vec{\boldsymbol{F}}_{\rm rec}^{{\rm x}(l)}\rangle \tag{2b}$$

$$\langle\vec{\boldsymbol{F}}_{\rm rec}^{{\rm e}(l)}\rangle = -\frac{1}{4\pi\varepsilon_0}\,\frac{(l+2)\,k^{2l+3}}{(l+1)!\,(2l+3)!!}\,{\rm Im}\Big[\overleftrightarrow{\mathbb{O}}_{\rm elec}^{(l)\,*}\overset{(l)}{\cdot\cdot}\overleftrightarrow{\mathbb{O}}_{\rm elec}^{(l+1)}\Big] \tag{2c}$$

$$\langle\vec{\boldsymbol{F}}_{\rm rec}^{{\rm m}(l)}\rangle = -\frac{\mu_0}{4\pi}\,\frac{(l+2)\,k^{2l+3}}{(l+1)!\,(2l+3)!!}\,{\rm Im}\Big[\overleftrightarrow{\mathbb{O}}_{\rm mag}^{(l)\,*}\overset{(l)}{\cdot\cdot}\overleftrightarrow{\mathbb{O}}_{\rm mag}^{(l+1)}\Big] \tag{2d}$$

$$\langle\vec{\boldsymbol{F}}_{\rm rec}^{{\rm x}(l)}\rangle = \frac{Z_0}{4\pi}\,\frac{k^{2l+2}}{l\,l!\,(2l+1)!!}\,{\rm Re}\big[\overleftrightarrow{\mathbb{O}}_{\rm elec}^{(l)}\overset{(l-1)}{\cdot\cdot}\overleftrightarrow{\mathbb{O}}_{\rm mag}^{(l)\,*}\big]\overset{(2)}{\cdot\cdot}\overleftrightarrow{\boldsymbol{\epsilon}}, \tag{2e}$$

where k is the wave number in the transparent (lossless) background medium where the particle is located, the quantities $\vec{\boldsymbol{E}}$ and $\vec{\boldsymbol{B}}$ denote, respectively, the incident electric and magnetic fields where the particle is immersed, $\overleftrightarrow{\boldsymbol{\epsilon}}$ is the Levi-Civita tensor, whose components ϵ_{ijk} are antisymmetric with respect to the permutation of any pair of indices, and the superscripts "e", "m", and "x" denote, respectively, the contribution due to the electric multipoles, magnetic multipoles and hybrid term. The electric and magnetic multipoles of order l are described by totally symmetric and traceless [6] rank-l tensors $\overleftrightarrow{\mathbb{O}}^{(l)}_{\rm elec}$ and $\overleftrightarrow{\mathbb{O}}^{(l)}_{\rm mag}$, respectively. The lower order cases with $l = 1$, 2, 3, 4, and 5 correspond to the dipole, quadrupole, octupole, hexadecapole, and dotriacontapole, respectively. The multiple contraction between two tensors of ranks l and l', denoted by $\overset{(m)}{\cdot\cdot}$, yields a tensor of rank $l + l' - 2m$ given by

$$\overleftrightarrow{\mathbb{A}}^{(l)} \overset{(m)}{\cdot\cdot} \overleftrightarrow{\mathbb{B}}^{(l')} = \mathbb{A}^{(l)}_{i_1\, i_2\, \cdots\, i_{l-m}\, k_1\, k_2\, \cdots\, k_{m-1}\, k_m} \mathbb{B}^{(l')}_{k_m\, k_{m-1}\, \cdots\, k_2\, k_1\, j_{m+1}\, \cdots\, j_{l'-1}\, j_{l'}}, \quad 0 \le m \le \min\left[\, l, l' \,\right], \tag{3}$$

with the summation over repeated indices assumed. Here the tensor contraction is made consecutively over two nearest indices in two index sequences. Equations (1) and (2) imply that the interception force is simply the coupling of the electric multipoles with the external electric field, governed by Eq. (1c), plus the coupling between the magnetic multipoles and the external magnetic field, determined by Eq. (1d). The recoil force originates from the coupling between the electric multipoles of adjacent orders, denoted by Eq. (2c), the magnetic multipoles of adjacent orders, delineated by Eq. (2d), and the electric and magnetic multipoles of the same order, depicted by Eq. (2e). Equations (1) and (2) constitute the complete multipole expansion of optical force on any particle, up to arbitrary orders of multipoles.

In the following, we give, for a spherical particle, the electric (magnetic) 2^l-pole moments, $\overleftrightarrow{\mathbb{O}}^{(l)}_{\rm elec}$ [$\overleftrightarrow{\mathbb{O}}^{(l)}_{\rm mag}$], in terms of the Mie coefficients a_l (b_l) [1] and the multiple gradient of the incident electric (magnetic) field. For a general (non-spherical) particle, the multipole moments are determined by its T-matrix [2], instead of the simple Mie coefficients. The T-matrix for a general non-spherical scatterer can only be evaluated numerically.

Define two totally symmetric tensors of rank n by

$$\begin{aligned}
\overleftrightarrow{\mathbb{M}}^{(n)}_{\rm elec} &= \hat{\mathcal{S}}\left[\, \nabla^{(n-1)} \vec{\boldsymbol{E}} \,\right] = \hat{\mathcal{S}}\left[\, \overbrace{\nabla\, \nabla \cdots \nabla}^{n-1}\, \vec{\boldsymbol{E}} \,\right] = \frac{1}{n} \sum_{j=1}^{n} \partial_{i_1} \cdots \partial_{i_{j-1}} \partial_{i_n} \partial_{i_{j+1}} \cdots \partial_{i_{n-1}} E_{i_j} \\
\overleftrightarrow{\mathbb{M}}^{(n)}_{\rm mag} &= \hat{\mathcal{S}}\left[\, \nabla^{(n-1)} \vec{\boldsymbol{B}} \,\right] = \hat{\mathcal{S}}\left[\, \overbrace{\nabla\, \nabla \cdots \nabla}^{n-1}\, \vec{\boldsymbol{B}} \,\right] = \frac{1}{n} \sum_{j=1}^{n} \partial_{i_1} \cdots \partial_{i_{j-1}} \partial_{i_n} \partial_{i_{j+1}} \cdots \partial_{i_{n-1}} B_{i_j}
\end{aligned} \tag{4}$$

where $\nabla^{(n)}$ means taking n consecutive gradients on a scalar or vector fields, whereas $\hat{\mathcal{S}}$ denotes the symmetrizing operator, E_j (B_j) is the j-th Cartesian component of the electric (magnetic) field, and ∂_i represents the partial derivative $\dfrac{\partial}{\partial x_i}$ with respect to the i-th Cartesian coordinate. The tensors $\overleftrightarrow{\mathbb{M}}^{(n)}_{\rm elec\,(mag)}$ so defined are obviously totally symmetric (invariant under the permutation of any pair of indices), but they are not totally traceless (vanishing under the contraction of any pair of indices) [6]. To obtain totally traceless tensors for the electric and magnetic multipoles, we proceed to define another two (electric and magnetic) sets of totally symmetric rank-l tensors by

$$\overleftrightarrow{\mathbb{N}}^{(l,\, m)}_{\rm elec\,(mag)} = \hat{\mathcal{S}}\left[\, \overbrace{\overleftrightarrow{\mathbb{I}} \otimes \overleftrightarrow{\mathbb{I}} \otimes \cdots \overleftrightarrow{\mathbb{I}} \otimes}^{m}\, \overleftrightarrow{\mathbb{M}}^{(l-2m)}_{\rm elec\,(mag)} \,\right] \quad \text{with} \quad \overleftrightarrow{\mathbb{N}}^{(l,\, 0)}_{\rm elec\,(mag)} = \overleftrightarrow{\mathbb{M}}^{(l)}_{\rm elec\,(mag)}, \tag{5}$$

where $\overleftrightarrow{\mathbb{I}}$ denotes the unit dyad of dimension 3, and the symbol $\otimes$ represents the tensor product so that the term in the square brackets means m consecutive times of tensor product, resulting in a rank-l tensor independent of m. In component form, it reads

$$\overleftrightarrow{\mathbb{N}}^{(l,\, m)}_{i_1 i_2, \cdots i_l} = \hat{\mathcal{S}}\left[\, \delta_{i_1 i_2} \delta_{i_3 i_4} \cdots \delta_{i_{l-2m-2} i_{l-2m-1}} \overleftrightarrow{\mathbb{M}}^{(l-2m)}_{i_{l-2m+1} i_{l-2m+2}, \cdots i_l} \,\right].$$

It can be proved that

$$\vec{\boldsymbol{n}} \times \left[\, \vec{\boldsymbol{n}}^{(l-2m-1)} \overset{(l-2m-1)}{\cdot\cdot} \overleftrightarrow{\mathbb{M}}^{(l-2m)}_{\rm elec\,(mag)} \,\right] = \frac{l}{l-2m}\, \vec{\boldsymbol{n}} \times \left[\, \vec{\boldsymbol{n}}^{(l-1)} \overset{(l-1)}{\cdot\cdot} \overleftrightarrow{\mathbb{N}}^{(l,\, m)}_{\rm elec\,(mag)} \,\right], \tag{6}$$

where $\vec{\boldsymbol{n}} = \vec{\boldsymbol{r}}/r$ is the unit vector in radial direction, and $\vec{\boldsymbol{n}}^{(m)}$ denotes a rank-m tensor resulting from the tensor product of m vectors $\vec{\boldsymbol{n}}$. For instance, $\vec{\boldsymbol{n}}^{(0)} = 1$, $\vec{\boldsymbol{n}}^{(1)} = \vec{\boldsymbol{n}}$, $\vec{\boldsymbol{n}}^{(2)} = \vec{\boldsymbol{n}}\otimes\vec{\boldsymbol{n}}$, and $\vec{\boldsymbol{n}}^{(3)} = \vec{\boldsymbol{n}}\otimes\vec{\boldsymbol{n}}\otimes\vec{\boldsymbol{n}}$. The totally symmetric and traceless multipole moments, $\overleftrightarrow{\mathbb{O}}^{(l)}_{\rm elec}$ and $\overleftrightarrow{\mathbb{O}}^{(l)}_{\rm mag}$, which are therefore usually referred to as 2^l-pole moments, are derived based on theory of multipole fields [4]

$$\begin{aligned}
&\overleftrightarrow{\mathbb{O}}^{(l)}_{\rm elec\,(mag)} = \gamma^{(l)}_{\rm elec\,(mag)} \sum_{m=0}^{\lfloor\frac{l-1}{2}\rfloor} d_{l,m}\, k^{2m}\, \overleftrightarrow{\mathbb{N}}^{(l,\,m)}_{\rm elec\,(mag)},\\
&d_{l,m} = \frac{1}{4^m}\,\frac{l!}{m!}\,\frac{\Gamma(l-m+\frac{1}{2})}{\Gamma(l+\frac{1}{2})\,\Gamma(l-2m)}\,\frac{1}{l}, \qquad \text{with} \quad d_{l,0} = 1,
\end{aligned} \tag{7}$$

where $\lfloor x\rfloor$ gives the greatest integer less than or equal to x and $\Gamma(x)$ denotes the Gamma function. The lower order cases with $l = 1$, 2, 3, 4, and 5 correspond to the dipole, quadrupole, octupole, hexadecapole, and dotriacontapole moments, respectively. For instance, $\overleftrightarrow{\mathbb{O}}^{(1)}_{\rm elec}$ reduces to the electric dipole moment $\vec{\boldsymbol{p}}$, $\overleftrightarrow{\mathbb{O}}^{(1)}_{\rm mag}$ is the magnetic dipole moment $\vec{\boldsymbol{m}}$, $\overleftrightarrow{\mathbb{O}}^{(2)}_{\rm elec}$ represents the electric quadrupole moment $\overleftrightarrow{\boldsymbol{Q}}^{(\boldsymbol{e})}$, and $\overleftrightarrow{\mathbb{O}}^{(2)}_{\rm mag}$ delineates the magnetic quadrupole moment $\overleftrightarrow{\boldsymbol{Q}}^{(\boldsymbol{m})}$, just to list a few lowest order cases. The electric and magnetic polarizabilities, $\gamma^{(l)}_{\rm elec}$ and $\gamma^{(l)}_{\rm mag}$, depend on the Mie coefficients [1] a_l and b_l of a spherical particle through

$$\gamma^{(l)}_{\rm elec} = \frac{4l(2l+1)!!}{(l+1)}\,\frac{i\pi\varepsilon_0\, a_l}{k^{2l+1}}, \qquad \gamma^{(l)}_{\rm mag} = \frac{4l(2l+1)!!}{(l+1)}\,\frac{i\pi\, b_l}{\mu_0 k^{2l+1}}. \tag{8}$$

In the Systèm International d'Unités (SI), the dimensions of $\gamma^{(l)}_{\rm mag}$ and $\gamma^{(l)}_{\rm elec}$ differ by a factor of c^2, viz, $\gamma^{(l)}_{\rm mag}/\gamma^{(l)}_{\rm elec} = c^2\,(b_l/a_l)$.

II. DECOMPOSITION INTO GRADIENT AND CURL FORCES

To decompose the optical force acting on a spherical particle into a sum of an irrotational term of zero curl and a solenoidal (divergenceless) term of zero divergence, or, equivalently, the conservative and non-conservative parts, it is convenient to write the optical force in terms of the multiple gradients $\overleftrightarrow{\mathbb{M}}^{(n)}_{\rm elec}$ and $\overleftrightarrow{\mathbb{M}}^{(n)}_{\rm mag}$ of the electric and magnetic fields defined in (4), instead of multipole moments $\overleftrightarrow{\mathbb{O}}^{(l)}_{\rm elec}$ and $\overleftrightarrow{\mathbb{O}}^{(l)}_{\rm mag}$. After lengthy algebra, the results read

$$\langle \vec{\boldsymbol{F}}^{\,\rm e(l)}_{\rm int}\rangle = \frac{1}{2\,l!}\,\mathrm{Re}\sum_{m=0}^{\lfloor\frac{l-1}{2}\rfloor} c_{l,m} k^{4m} \gamma^{(l)}_{\rm elec}\big[\nabla^{(l-2m)}\vec{\boldsymbol{E}}^{*}\big]\,\overset{(l-2m)}{\cdot\cdot}\,\overleftrightarrow{\mathbb{M}}^{(l\,-\,2m)}_{\rm elec}, \tag{9a}$$

$$\langle \vec{\boldsymbol{F}}^{\,\rm m(l)}_{\rm int}\rangle = \frac{1}{2\,l!}\,\mathrm{Re}\sum_{m=0}^{\lfloor\frac{l-1}{2}\rfloor} c_{l,m} k^{4m} \gamma^{(l)}_{\rm mag}\big[\nabla^{(l-2m)}\vec{\boldsymbol{B}}^{*}\big]\,\overset{(l-2m)}{\cdot\cdot}\,\overleftrightarrow{\mathbb{M}}^{(l\,-\,2m)}_{\rm mag}, \tag{9b}$$

$$\begin{aligned}
\langle \vec{\boldsymbol{F}}^{\,\rm e(l)}_{\rm rec}\rangle = &-\frac{1}{4\pi\varepsilon_0}\,\frac{(l+2)\,k^{2l+3}}{(l+1)!\,(2l+3)!!}\,\mathrm{Im}\sum_{m=0}^{\lfloor\frac{l-2}{2}\rfloor} g_{l,m}\,k^{4m+2}\,\eta^{(l)}_{\rm elec}\,\overleftrightarrow{\mathbb{M}}^{(l\,-\,2m)\,*}_{\rm elec}\,\overset{(l-2m-1)}{\cdot\cdot}\,\overleftrightarrow{\mathbb{M}}^{(l\,-\,2m\,-\,1)}_{\rm elec}\\
&-\frac{1}{4\pi\varepsilon_0}\,\frac{(l+2)\,k^{2l+3}}{(l+1)!\,(2l+3)!!}\,\mathrm{Im}\sum_{m=0}^{\lfloor\frac{l-1}{2}\rfloor} f_{l,m}\,k^{4m}\,\eta^{(l)}_{\rm elec}\,\overleftrightarrow{\mathbb{M}}^{(l\,-\,2m)\,*}_{\rm elec}\,\overset{(l-2m)}{\cdot\cdot}\,\overleftrightarrow{\mathbb{M}}^{(l\,-\,2m\,+\,1)}_{\rm elec},
\end{aligned} \tag{10a}$$

$$\begin{aligned}
\langle \vec{\boldsymbol{F}}^{\,\rm m(l)}_{\rm rec}\rangle = &-\frac{\mu_0}{4\pi}\,\frac{(l+2)\,k^{2l+3}}{(l+1)!\,(2l+3)!!}\,\mathrm{Im}\sum_{m=0}^{\lfloor\frac{l-2}{2}\rfloor} g_{l,m}\,k^{4m+2}\,\eta^{(l)}_{\rm mag}\,\overleftrightarrow{\mathbb{M}}^{(l\,-\,2m)\,*}_{\rm mag}\,\overset{(l-2m-1)}{\cdot\cdot}\,\overleftrightarrow{\mathbb{M}}^{(l\,-\,2m\,-\,1)}_{\rm mag}\\
&-\frac{\mu_0}{4\pi}\,\frac{(l+2)\,k^{2l+3}}{(l+1)!\,(2l+3)!!}\,\mathrm{Im}\sum_{m=0}^{\lfloor\frac{l-1}{2}\rfloor} f_{l,m}\,k^{4m}\,\eta^{(l)}_{\rm mag}\,\overleftrightarrow{\mathbb{M}}^{(l\,-\,2m)\,*}_{\rm mag}\,\overset{(l-2m)}{\cdot\cdot}\,\overleftrightarrow{\mathbb{M}}^{(l\,-\,2m\,+\,1)}_{\rm mag},
\end{aligned} \tag{10b}$$

$$\langle \vec{\boldsymbol{F}}_{\rm rec}^{\,{\rm x}(l)} \rangle \;=\; \frac{Z_0}{4\pi}\,\frac{k^{2l+2}}{l\,l!\,(2l+1)!!}\,{\rm Re}\sum_{m=0}^{\lfloor\frac{l-1}{2}\rfloor} h_{l,m}\,k^{4m}\eta_{\rm hyb}^{(l)}\big[\,\overleftrightarrow{\mathbb{M}}_{\rm elec}^{(l-2m)}\;\overset{(l-2m-1)}{\cdot\cdot}\;\overleftrightarrow{\mathbb{M}}_{\rm mag}^{(l-2m)\,*}\,\big]\overset{(2)}{\cdot\cdot}\overleftrightarrow{\boldsymbol{\epsilon}}, \tag{10c}$$

where $\eta_{\rm elec}^{(l)} = \gamma_{\rm elec}^{(l+1)}\gamma_{\rm elec}^{(l)\,*}$, $\eta_{\rm mag}^{(l)} = \gamma_{\rm mag}^{(l+1)}\gamma_{\rm mag}^{(l)\,*}$, and $\eta_{\rm hyb}^{(l)} = \gamma_{\rm elec}^{(l)}\gamma_{\rm mag}^{(l)\,*}$, are products of polarizabilities. In deriving Eqs. (9) and (10), we have used the following mathematical identities

$$\begin{aligned}\overleftrightarrow{\mathbb{O}}_{\rm elec\,(mag)}^{(l)\,*}\;\overset{(l)}{\cdot\cdot}\;\overleftrightarrow{\mathbb{O}}_{\rm elec\,(mag)}^{(l+1)} \;=\; &\sum_{m=0}^{\lfloor(l-1)/2\rfloor} f_{l,m}\,k^{4m}\,\eta_{\rm elec\,(mag)}^{(l)}\,\overleftrightarrow{\mathbb{M}}_{\rm elec\,(mag)}^{(l-2m)\,*}\;\overset{(l-2m)}{\cdot\cdot}\;\overleftrightarrow{\mathbb{M}}_{\rm elec\,(mag)}^{(l-2m+1)}\\ &+\sum_{m=0}^{\lfloor(l-2)/2\rfloor} g_{l,m}\,k^{4m+2}\,\eta_{\rm elec\,(mag)}^{(l)}\,\overleftrightarrow{\mathbb{M}}_{\rm elec\,(mag)}^{(l-2m)\,*}\;\overset{(l-2m-1)}{\cdot\cdot}\;\overleftrightarrow{\mathbb{M}}_{\rm elec\,(mag)}^{(l-2m-1)},\end{aligned} \tag{11a}$$

and

$$\big[\,\overleftrightarrow{\mathbb{O}}_{\rm elec}^{(l)}\;\overset{(l-1)}{\cdot\cdot}\;\overleftrightarrow{\mathbb{O}}_{\rm mag}^{(l)\,*}\,\big]\overset{(2)}{\cdot\cdot}\overleftrightarrow{\boldsymbol{\epsilon}} = \sum_{m=0}^{\lfloor(l-1)/2\rfloor} h_{l,m}\,k^{4m}\big[\,\eta_{\rm hyb}^{(l)}\,\overleftrightarrow{\mathbb{M}}_{\rm elec}^{(l-2m)}\;\overset{(l-2m-1)}{\cdot\cdot}\;\overleftrightarrow{\mathbb{M}}_{\rm mag}^{(l-2m)\,*}\,\big]\overset{(2)}{\cdot\cdot}\overleftrightarrow{\boldsymbol{\epsilon}} \tag{11b}$$

with

$$c_{l,m} \;=\; \frac{(-1)^m}{4^m}\,\frac{l!}{m!}\,\frac{\Gamma(l-m+\frac{1}{2})}{\Gamma(l+\frac{1}{2})\,\Gamma(l-2m)}\,\frac{(l-2m)}{l^2} = \frac{(-1)^m(l-2m)}{l}\,d_{l,m}, \tag{11c}$$

$$f_{l,m} \;=\; \frac{(-1)^m}{4^m}\,\frac{l!}{m!}\,\frac{\Gamma(l-m+\frac{1}{2})}{\Gamma(l+\frac{1}{2})\,\Gamma(l-2m)}\,\frac{(l-2m+1)(2l-2m+1)}{l(l+1)(2l+1)}, \tag{11d}$$

$$g_{l,m} \;=\; \frac{(-1)^m}{4^m}\,\frac{l!}{m!}\,\frac{\Gamma(l-m+\frac{1}{2})}{\Gamma(l+\frac{1}{2})\,\Gamma(l-2m)}\,\frac{(l-2m)(l-2m-1)}{l(l+1)(2l+1)}, \tag{11e}$$

$$h_{l,m} \;=\; \frac{(-1)^m}{4^m}\,\frac{l!}{m!}\,\frac{\Gamma(l-m+\frac{1}{2})}{\Gamma(l+\frac{1}{2})\,\Gamma(l-2m)}\,\frac{(l-2m)^2}{l^3}. \tag{11f}$$

The decomposition of the optical force into the irrotational and the solenoidal terms starts with Eqs. (9) and (10) as well as the definitions Eq. (4). To achieve high readability, we assume the particle is located in the regime with $z > 0$ and write the incident wave, in which the particle is immersed, in terms of the plane wave spectrum representation [7–12], given below,

$$\vec{\boldsymbol{E}} = \vec{\boldsymbol{E}}_{\rm inc} = \iint_{-\infty}^{\infty} \vec{\boldsymbol{e}}_{\vec{\boldsymbol{u}}}\,e^{ik\vec{\boldsymbol{u}}\cdot\vec{\boldsymbol{r}}}\,{\rm d}u_x\,{\rm d}u_y \quad\text{and}\quad \vec{\boldsymbol{H}} = \vec{\boldsymbol{H}}_{\rm inc} = \frac{1}{Z_0}\iint_{-\infty}^{\infty} \vec{\boldsymbol{h}}_{\vec{\boldsymbol{u}}}\,e^{ik\vec{\boldsymbol{u}}\cdot\vec{\boldsymbol{r}}}\,{\rm d}u_x\,{\rm d}u_y, \tag{12}$$

where k is the wave number in the background medium,

$$\vec{\boldsymbol{u}} = u_x\,\vec{\boldsymbol{e}}_x + u_y\,\vec{\boldsymbol{e}}_y + u_z\,\vec{\boldsymbol{e}}_z, \quad \text{with}\;\; u_z = \begin{cases}\sqrt{1-u_x^2-u_y^2}, & \text{if } u_x^2+u_y^2 \le 1\\ i\sqrt{u_x^2+u_y^2-1}, & \text{if } u_x^2+u_y^2 > 1\end{cases} \quad\text{and}\quad \vec{\boldsymbol{u}}\cdot\vec{\boldsymbol{u}} = 1,$$

whereas the electric and magnetic plane wave spectra $\vec{\boldsymbol{e}}_{\vec{\boldsymbol{u}}}$ and $\vec{\boldsymbol{h}}_{\vec{\boldsymbol{u}}}$ satisfy

$$\vec{\boldsymbol{u}}\cdot\vec{\boldsymbol{e}}_{\vec{\boldsymbol{u}}} = \vec{\boldsymbol{u}}\cdot\vec{\boldsymbol{h}}_{\vec{\boldsymbol{u}}} = 0, \quad \vec{\boldsymbol{h}}_{\vec{\boldsymbol{u}}} = \vec{\boldsymbol{u}}\times\vec{\boldsymbol{e}}_{\vec{\boldsymbol{u}}}, \quad \vec{\boldsymbol{e}}_{\vec{\boldsymbol{u}}} = -\vec{\boldsymbol{u}}\times\vec{\boldsymbol{h}}_{\vec{\boldsymbol{u}}}. \tag{13}$$

With the plane wave spectrum representation Eqs. (12), the multiple gradients in (4) can be written as

$$\overleftrightarrow{\mathbb{M}}_{\rm elec}^{(n)} \;=\; \frac{(ik)^{n-1}}{n}\sum_{j=0}^{n-1}\iint_{-\infty}^{\infty}\vec{\boldsymbol{u}}^{(n-1-j)}\;\vec{\boldsymbol{e}}_{\vec{\boldsymbol{u}}}\,\vec{\boldsymbol{u}}^{(j)}\,e^{ik\vec{\boldsymbol{u}}\cdot\vec{\boldsymbol{r}}}\,{\rm d}u_x\,{\rm d}u_y, \tag{14a}$$

$$\overleftrightarrow{\mathbb{M}}_{\rm mag}^{(n)} \;=\; \frac{(ik)^{n-1}}{n\,c}\sum_{j=0}^{n-1}\iint_{-\infty}^{\infty}\vec{\boldsymbol{u}}^{(n-1-j)}\;\vec{\boldsymbol{h}}_{\vec{\boldsymbol{u}}}\,\vec{\boldsymbol{u}}^{(j)}\,e^{ik\vec{\boldsymbol{u}}\cdot\vec{\boldsymbol{r}}}\,{\rm d}u_x\,{\rm d}u_y, \tag{14b}$$

where $\vec{\boldsymbol{u}}^{(n)}$ denotes the tensor product of n vectors $\vec{\boldsymbol{u}}$, viz, $\vec{\boldsymbol{u}}^{(0)}=1$, $\vec{\boldsymbol{u}}^{(1)}=\vec{\boldsymbol{u}}$, $\vec{\boldsymbol{u}}^{(2)}=\vec{\boldsymbol{u}}\otimes\vec{\boldsymbol{u}}$, and $\vec{\boldsymbol{u}}^{(3)}=\vec{\boldsymbol{u}}\otimes\vec{\boldsymbol{u}}\otimes\vec{\boldsymbol{u}}$, with the symbol $\otimes$ representing the tensor product.

After lengthy algebra, the interception parts $\langle\vec{\boldsymbol{F}}_{\rm int}^{\rm e(}{}^{l)}\rangle$ and $\langle\vec{\boldsymbol{F}}_{\rm int}^{\rm m(}{}^{l)}\rangle$ of the optical force involving order l multipoles can be rewritten as

$$\begin{aligned}\langle\vec{\boldsymbol{F}}_{\rm int}^{\rm e(}{}^{l)}\rangle &= \frac{1}{2\,l!}\sum_{m=0}^{\lfloor\frac{l-1}{2}\rfloor} c_{lm}k^{4m}\,{\rm Re}\left[\,\gamma_{\rm elec}^{(l)}\,\vec{\boldsymbol{t}}_{\rm elec}^{(l-2m)}\,\right]\\ \langle\vec{\boldsymbol{F}}_{\rm int}^{\rm m(}{}^{l)}\rangle &= \frac{1}{2\,l!}\sum_{m=0}^{\lfloor\frac{l-1}{2}\rfloor} c_{lm}k^{4m}\,{\rm Re}\left[\,\gamma_{\rm mag}^{(l)}\,\vec{\boldsymbol{t}}_{\rm mag}^{(l-2m)}\,\right]\end{aligned}\tag{15a}$$

where

$$\begin{aligned}\vec{\boldsymbol{t}}_{\rm elec}^{(n)} &\equiv \left[\nabla^{(n)}\vec{\boldsymbol{E}}^{*}\right]\,\vdots\!\vdots^{(n)}\,\overleftrightarrow{\mathbb{M}}_{\rm elec}^{(n)}\\ &= \frac{1}{2}\left[\nabla D_{\rm ee}^{(n)}-\nabla\times\vec{\boldsymbol{S}}_{\rm ee}^{(n)}-2i\omega\,{\rm Re}\,\vec{\boldsymbol{S}}_{\rm em}^{(n)}\right]\\ &\quad-\frac{(n-1)\omega^2}{2n}\left[\nabla D_{\rm mm}^{(n-1)}-\nabla\times\vec{\boldsymbol{S}}_{\rm mm}^{(n-1)}-\frac{2i\omega}{c^2}\,{\rm Re}\,\vec{\boldsymbol{S}}_{\rm em}^{(n-1)}\right],\end{aligned}\tag{15b}$$

$$\begin{aligned}\vec{\boldsymbol{t}}_{\rm mag}^{(n)} &\equiv \left[\nabla^{(n)}\vec{\boldsymbol{B}}^{*}\right]\,\vdots\!\vdots^{(n)}\,\overleftrightarrow{\mathbb{M}}_{\rm mag}^{(n)}\\ &= \frac{1}{2}\left[\nabla D_{\rm mm}^{(n)}-\nabla\times\vec{\boldsymbol{S}}_{\rm mm}^{(n)}-\frac{2i\omega}{c^2}\,{\rm Re}\,\vec{\boldsymbol{S}}_{\rm em}^{(n)}\right]\\ &\quad-\frac{(n-1)\omega^2}{2n\,c^4}\left[\nabla D_{\rm ee}^{(n-1)}-\nabla\times\vec{\boldsymbol{S}}_{\rm ee}^{(n-1)}-2i\omega\,{\rm Re}\,\vec{\boldsymbol{S}}_{\rm em}^{(n-1)}\right].\end{aligned}\tag{15c}$$

The field moments in the reciprocal space are defined as follows.

$$\begin{aligned}D_{\rm ee}^{(n)} &= \left(\nabla^{(n-1)}\vec{\boldsymbol{E}}\right)\,\vdots^{(n)}\,\left(\nabla^{(n-1)}\vec{\boldsymbol{E}}^{*}\right),\\ D_{\rm mm}^{(n)} &= \left(\nabla^{(n-1)}\vec{\boldsymbol{B}}\right)\,\vdots^{(n)}\,\left(\nabla^{(n-1)}\vec{\boldsymbol{B}}^{*}\right),\\ \vec{\boldsymbol{S}}_{\rm ee}^{(n)} &= \left[\left(\nabla^{(n-1)}\vec{\boldsymbol{E}}\right)\,\vdots^{(n-1)}\,\left(\nabla^{(n-1)}\vec{\boldsymbol{E}}^{*}\right)\right]\,\vdots^{(2)}\,\overleftrightarrow{\boldsymbol{\epsilon}},\\ \vec{\boldsymbol{S}}_{\rm mm}^{(n)} &= \left[\left(\nabla^{(n-1)}\vec{\boldsymbol{B}}\right)\,\vdots^{(n-1)}\,\left(\nabla^{(n-1)}\vec{\boldsymbol{B}}^{*}\right)\right]\,\vdots^{(2)}\,\overleftrightarrow{\boldsymbol{\epsilon}},\\ \vec{\boldsymbol{S}}_{\rm em}^{(n)} &= \left[\left(\nabla^{(n-1)}\vec{\boldsymbol{E}}\right)\,\vdots^{(n-1)}\,\left(\nabla^{(n-1)}\vec{\boldsymbol{B}}^{*}\right)\right]\,\vdots^{(2)}\,\overleftrightarrow{\boldsymbol{\epsilon}}.\end{aligned}\tag{16}$$

It is noted that $D_{\rm ee}^{(n)}$ and $D_{\rm mm}^{(n)}$ are real while $\vec{\boldsymbol{S}}_{\rm ee}^{(n)}$ and $\vec{\boldsymbol{S}}_{\rm mm}^{(n)}$ are purely imaginary. In Eqs. (16) we have defined the second kind of multiple tensor contraction, denoted by $\vdots^{(m)}$,

$$\overleftrightarrow{\mathbb{A}}^{(n)}\,\vdots^{(m)}\,\overleftrightarrow{\mathbb{B}}^{(n')}=\mathbb{A}^{(n)}_{k_1\,k_2\,\cdots\,k_m\,i_{m+1}\,i_{m+2}\,\cdots\,i_n}\,\mathbb{B}^{(n')}_{k_1\,k_2\,\cdots\,k_m\,j_{m+1}\,j_{m+2}\,\cdots\,j_{n'}},\qquad 0\le m\le\min[n,n']\tag{17}$$

which differs from Eq. (3) in that the tensor contraction is made over the corresponding left most indices, instead of over the nearest indices, in the two index sequences. Some simple examples are

$$\begin{aligned}&(\vec{\boldsymbol{v}}\vec{\boldsymbol{w}})\,\vdots\!\vdots^{(2)}\,\overleftrightarrow{\boldsymbol{\epsilon}}=\overleftrightarrow{\boldsymbol{\epsilon}}\,\vdots\!\vdots^{(2)}\,(\vec{\boldsymbol{v}}\vec{\boldsymbol{w}})=\vec{\boldsymbol{w}}\times\vec{\boldsymbol{v}},\qquad&&\text{versus}\qquad&&(\vec{\boldsymbol{v}}\vec{\boldsymbol{w}})\,\vdots^{(2)}\,\overleftrightarrow{\boldsymbol{\epsilon}}=\overleftrightarrow{\boldsymbol{\epsilon}}\,\vdots^{(2)}\,(\vec{\boldsymbol{v}}\vec{\boldsymbol{w}})=\vec{\boldsymbol{v}}\times\vec{\boldsymbol{w}},\\ &(\nabla\nabla\vec{\boldsymbol{E}})\,\vdots\!\vdots^{(2)}\,(\nabla\vec{\boldsymbol{B}})=(\partial_i\partial_j E_k)(\partial_k B_j)\qquad&&\text{versus}\qquad&&(\nabla\nabla\vec{\boldsymbol{E}})\,\vdots^{(2)}\,(\nabla\vec{\boldsymbol{B}})=(\partial_i\partial_j E_k)(\partial_i B_j),\\ &(\nabla\vec{\boldsymbol{B}})\,\vdots\!\vdots^{(2)}\,(\nabla\nabla\vec{\boldsymbol{E}})=(\partial_i B_j)(\partial_j\partial_i E_k)\qquad&&\text{versus}\qquad&&(\nabla\vec{\boldsymbol{B}})\,\vdots^{(2)}\,(\nabla\nabla\vec{\boldsymbol{E}})=(\partial_i B_j)(\partial_i\partial_j E_k).\end{aligned}\tag{18}$$

Since ${\rm Re}\,\vec{\boldsymbol{S}}_{\rm em}^{(n)}$ is solenoidal (divergenceless, as can be seen below), Eq. (15), together with (1a) and (1b), concludes the decomposition of the interception (extinction) force. In the dipolar limit, one has $l=1$, the interception force can be written in terms of gradients $\nabla D_{\rm ee}^{(1)}=\nabla|\vec{\boldsymbol{E}}|^2$ and $\nabla D_{\rm mm}^{(n)}=\nabla|\vec{\boldsymbol{B}}|^2$ of field intensity, the Poynting vector $\vec{\boldsymbol{S}}_{\rm em}^{(1)}=\vec{\boldsymbol{E}}\times\vec{\boldsymbol{B}}^{*}$, and the curl of spin angular momentum density, $\nabla\times(\vec{\boldsymbol{E}}\times\vec{\boldsymbol{E}}^{*})$ and $\nabla\times(\vec{\boldsymbol{B}}\times\vec{\boldsymbol{B}}^{*})$, which has been claimed by many authors as the origin of the optical force (see, e.g., Ref. [14]).

The electric and magnetic parts of the recoil force can be worked out to give

$$\begin{aligned}\langle \vec{\boldsymbol{F}}_{\rm rec}^{\,{\rm e}(l)} \rangle &= -\frac{c_l\, k^{2l+3}}{4\pi\varepsilon_0}\Big\{ \sum_{m=0}^{\lfloor (l-1)/2 \rfloor} f_{l,m} k^{4m}\, {\rm Im}\big[\eta_{\rm elec}^{(l)} \vec{\boldsymbol{t}}_{\rm ee}^{\,(l-2m)\,*}\big] + \sum_{m=0}^{\lfloor (l-2)/2 \rfloor} g_{l,m} k^{4m+2}\, {\rm Im}\big[\eta_{\rm elec}^{(l)} \vec{\boldsymbol{t}}_{\rm ee}^{\,(l-2m-1)}\big]\Big\} \\ \langle \vec{\boldsymbol{F}}_{\rm rec}^{\,{\rm m}(l)} \rangle &= -\frac{c_l\, k^{2l+3}}{4\pi\varepsilon_0 c^2}\Big\{ \sum_{m=0}^{\lfloor (l-1)/2 \rfloor} f_{l,m} k^{4m}\, {\rm Im}\big[\eta_{\rm mag}^{(l)} \vec{\boldsymbol{t}}_{\rm mm}^{\,(l-2m)\,*}\big] + \sum_{m=0}^{\lfloor (l-2)/2 \rfloor} g_{l,m} k^{4m+2}\, {\rm Im}\big[\eta_{\rm mag}^{(l)} \vec{\boldsymbol{t}}_{\rm mm}^{\,(l-2m-1)}\big]\Big\}\end{aligned} \tag{19}$$

where

$$\begin{aligned}\vec{\boldsymbol{t}}_{\rm ee}^{\,(n)} &\equiv \overleftrightarrow{\mathbb{M}}_{\rm elec}^{(n)} \overset{(n)}{\cdot\cdot} \overleftrightarrow{\mathbb{M}}_{\rm elec}^{(n+1)\,*} = \overleftrightarrow{\mathbb{M}}_{\rm elec}^{(n+1)\,*} \overset{(n)}{\cdot\cdot} \overleftrightarrow{\mathbb{M}}_{\rm elec}^{(n)} \\ &= \frac{1}{2}\Big[\nabla D_{\rm ee}^{(n)} - \nabla\times \vec{\boldsymbol{S}}_{\rm ee}^{(n)} - 2i\omega\, {\rm Re}\, \vec{\boldsymbol{S}}_{\rm em}^{(n)}\Big] + \frac{i\omega}{(n+1)}\, S_{\rm em}^{(n)} \\ &\quad - \frac{(n-1)\omega^2}{2(n+1)}\Big[\nabla D_{\rm mm}^{(n-1)} - \nabla\times \vec{\boldsymbol{S}}_{\rm mm}^{(n-1)} - \frac{2i\omega}{c^2}\, {\rm Re}\, \vec{\boldsymbol{S}}_{\rm em}^{(n-1)}\Big]\end{aligned} \tag{20a}$$

$$\begin{aligned}\vec{\boldsymbol{t}}_{\rm mm}^{\,(n)} &\equiv \overleftrightarrow{\mathbb{M}}_{\rm mag}^{(n)} \overset{(n)}{\cdot\cdot} \overleftrightarrow{\mathbb{M}}_{\rm mag}^{(n+1)\,*} = \overleftrightarrow{\mathbb{M}}_{\rm mag}^{(n+1)\,*} \overset{(n)}{\cdot\cdot} \overleftrightarrow{\mathbb{M}}_{\rm mag}^{(n)} \\ &= \frac{1}{2}\Big[\nabla D_{\rm mm}^{(n)} - \nabla\times \vec{\boldsymbol{S}}_{\rm mm}^{(n)} - \frac{2i\omega}{c^2}\, {\rm Re}\, \vec{\boldsymbol{S}}_{\rm em}^{(n)}\Big] + \frac{i\omega}{(n+1)c^2}\, S_{\rm em}^{(n)\,*} \\ &\quad - \frac{(n-1)\omega^2}{2(n+1)c^4}\Big[\nabla D_{\rm ee}^{(n-1)} - \nabla\times \vec{\boldsymbol{S}}_{\rm ee}^{(n-1)} - 2i\omega\, {\rm Re}\, \vec{\boldsymbol{S}}_{\rm em}^{(n-1)}\Big]\end{aligned} \tag{20b}$$

with $c_l = (l+2)/[(l+1)!(2l+3)!!]$.

Finally, the hybrid term $\langle \vec{\boldsymbol{F}}_{\rm rec}^{\,{\rm x}(l)} \rangle$ of the recoil force, given by (10c), can be cast, for our purpose, into

$$\langle \vec{\boldsymbol{F}}_{\rm rec}^{\,{\rm x}(l)} \rangle = \frac{Z_0}{4\pi}\, \frac{k^{2l+2}}{l\, l!(2l+1)!!} \sum_{m=0}^{\lfloor (l-1)/2 \rfloor} h_{l,m} k^{4m}\, {\rm Re}\big[\eta_{\rm hyb}^{(l)} \vec{\boldsymbol{t}}_{\rm em}^{\,(l-2m)}\big]. \tag{21}$$

where, written in a symmetric form with respect to the electric and magnetic fields (possessing so called "electric-magnetic democracy" [13]),

$$\begin{aligned}\vec{\boldsymbol{t}}_{\rm em}^{\,(n)} &\equiv \big[\overleftrightarrow{\mathbb{M}}_{\rm elec}^{(n)} \overset{(n-1)}{\cdot\cdot} \overleftrightarrow{\mathbb{M}}_{\rm mag}^{(n)\,*}\big] \overset{(2)}{\cdot\cdot} \overleftrightarrow{\boldsymbol{\epsilon}} \\ &= -\frac{i(n-1)\omega}{2n\, c^2}\nabla\times\big[\vec{\boldsymbol{S}}_{\rm ee}^{(n-1)} + c^2 \vec{\boldsymbol{S}}_{\rm mm}^{(n-1)}\big] + \frac{i(n-1)(n-2)\omega k^2}{2n^2 c^2}\nabla\times\big[\vec{\boldsymbol{S}}_{\rm ee}^{(n-2)} + c^2 \vec{\boldsymbol{S}}_{\rm mm}^{(n-2)}\big] - \frac{1}{n}\,\vec{\boldsymbol{S}}_{\rm em}^{(n)} \\ &\quad - \frac{(n-1)}{2n}\nabla\times\big[\vec{\boldsymbol{G}}_{\rm em}^{(n-1)} - \vec{\boldsymbol{G}}_{\rm me}^{(n-1)\,*}\big] + \frac{(n-1)(n-2)k^2}{2n^2}\nabla\times\big[\vec{\boldsymbol{G}}_{\rm em}^{(n-2)\,*} - \vec{\boldsymbol{G}}_{\rm me}^{(n-2)}\big] + \frac{(n-1)k^2}{n^2}\,\vec{\boldsymbol{S}}_{\rm em}^{(n-1)\,*}\end{aligned} \tag{22}$$

The two new field moments, $\vec{\boldsymbol{G}}_{\rm em}^{(n)}$ and $\vec{\boldsymbol{G}}_{\rm me}^{(n)}$, in the reciprocal space are given by

$$\vec{\boldsymbol{G}}_{\rm em}^{(n)} = [\nabla^{(n-1)}\vec{\boldsymbol{E}}] \overset{(n)}{\vdots} [\nabla^{(n)}\vec{\boldsymbol{B}}^*], \qquad \vec{\boldsymbol{G}}_{\rm me}^{(n)} = [\nabla^{(n-1)}\vec{\boldsymbol{B}}] \overset{(n)}{\vdots} [\nabla^{(n)}\vec{\boldsymbol{E}}^*]. \tag{23}$$

In the dipolar limit, with $l = 1$, the hybrid term depends only on the complex Poynting vector $\vec{\boldsymbol{S}}_{\rm em}^{(1)} = \vec{\boldsymbol{E}}\times\vec{\boldsymbol{B}}^*$, which is interpreted as a lateral optical force resulting from the Belinfante spin momentum [15, 16] in two-wave interference.

The recoil force, given by (19 - 22), depends on $\vec{\boldsymbol{S}}_{\rm em}^{(n)}$, which is neither irrotational nor solenoidal. So we need to decompose $\vec{\boldsymbol{S}}_{\rm em}^{(n)}$ into a gradient and curl parts. To this end, we write

$$\vec{\boldsymbol{S}}_{\rm em}^{(n)} = -\nabla\varphi_s^{(n)} + \nabla\times\vec{\boldsymbol{\psi}}_s^{(n)}. \tag{24}$$

It is worked out that

$$\varphi_s^{(n)} = -\frac{ik^{2n}}{2\omega}\sum_{m=n}^{\infty}\frac{1}{k^{2m}}\big[D_{\rm ee}^{(m)} - c^2 D_{\rm mm}^{(m)}\big], \tag{25a}$$

$$\vec{\boldsymbol{\psi}}_s^{(n)} = -\frac{k^{2(n-1)}}{2}\sum_{m=n}^{\infty}\frac{1}{k^{2m}}\Big[\vec{\boldsymbol{G}}_{\rm em}^{(m)} - \vec{\boldsymbol{G}}_{\rm me}^{(m)*}\Big], \tag{25b}$$

which result in

$$\vec{\boldsymbol{S}}^{(n)}_{\rm em} = -\frac{k^{2n-2}}{2}\nabla\times\sum_{m=n}^{\infty}\frac{1}{k^{2m}}\big[\vec{\boldsymbol{G}}^{(m)}_{\rm em} - \vec{\boldsymbol{G}}^{(m)*}_{\rm me}\big] + \frac{ik^{2n}}{2\omega}\nabla\sum_{m=n}^{\infty}\frac{1}{k^{2m}}\big[D^{(m)}_{\rm ee} - c^2 D^{(m)}_{\rm mm}\big]. \tag{26}$$

Since $D^{(n)}_{\rm ee}$ and $D^{(n)}_{\rm mm}$ are real, the real part $\mathrm{Re}\,\vec{\boldsymbol{S}}^{(n)}_{\rm em}$ is divergenceless, as stated in the text after Eq. (18). Equations (19 - 22), together with (26), (2a), and (2b), constitute the decomposition of the recoil force into a conservative and a nonconservative parts.

[1] C. F. Bohren and D. R. Huffman, *Absorption and Scattering of Light by Small Particles*, (JohnWiley, New York, 1998).

[2] M. I. Mishchenko, N. T. Zakharov, N. G. Khlebtsov, T. Wriedt, and G. Videen, J. Quant. Spectrosc. Radiat. Transfer. 146, 349 (2014) and references therein.

[3] J. A. Stratton, *Electromagnetic Theory*, (McGraw Hill, New York, 1941).

[4] M. E. Rose, *Multipole Fields*, (John Wiley, New York, 1951).

[5] B. L. Silver, *Irreducible Tensor Methods*, (Academic Press, New York, 1976).

[6] A totally symmetric tensor is a tensor with its components being symmetric with respect to the permutation of any pair of indices, while a totally traceless tensor is a tensor that vanishes under the contraction over any pair of indices. see, J. Applequist, J. Phys. A **22**, 4303 (1989).

[7] P. C. Clemmow, *The Plane Wave Spectrum Representation of Electromagnetic Fields* (Pergamon Press, Oxford, 1966).

[8] L. Mandel and E. Wolf, *Optical Coherence and Quantum Optics*, (Cambridge Univ. Press, Cambridge 1995).

[9] M. Nieto-Vesperinas, *Scattering and Diffraction in Physical Optics*, (World Scientific, Singapore 2006).

[10] J. W. Goodman, *Introduction to Fourier Optics* (Roberts & Company Publishers, Green Village 2005).

[11] S. Colak, C. Yeh, and L. W. Casperson, Appl. Opt. **18**, 294 (1979).

[12] G. Gouesbet and J. A. Lock, J. Quant. Spectrosc. Radiat. Transf. **162**, 18 (2015).

[13] M. V. Berry, J. Opt. A **11**, 094001 (2009).

[14] S. Albaladejo, M. I. Marques, M. Laroche, and J. J. Saenz, Phys. Rev. Lett. **102**, 113602 (2009).

[15] A. Y. Bekshaev, K. Y. Bliokh, and F. Nori, Phys. Rev. X **5**, 011039 (2015).

[16] F. J. Belinfante, Physica (Amsterdam) **7**, 449 (1940).